

CLINTAS CL

COMPOSITION : Each Vaginal Suppository contains :Clindamycin Phosphate USP equivalent to Clindamycin 100 mg and Clotrimazole IP 200 mg **Indication:** It is indicated for the treatment of mixed vaginitis or bacterial vaginosis coexisting with candidiasis, bacterial vaginitis and symptomatic treatment of mixed vaginal infections where *Trichomonas vaginalis* is also present. **DOSAGE & ADMINISTRATION:** One clintas CL vaginal suppository should be inserted as deeply as possible into the vagina, preferably on retiring to bed, for 3 consecutive nights. This is best achieved when lying on the back with legs pulled in a little towards the body. The treatment should not be carried out during menstruation but should be completed before onset of menstruation. The safety and efficacy in premenarchal females have not been established. **CONTRAINDICATIONS:** In patients with a history of hypersensitivity to clindamycin, lincomycin, clotrimazole or any other excipient of the formulation. In patients with a history of regional enteritis, ulcerative colitis or a history of "antibiotic associated" colitis. **Warnings & Precautions:** For vaginal use only. Not to be taken orally. Diarrhea, bloody diarrhea, and colitis (including pseudomembranous colitis) have been reported with the use of orally and parenterally administered clindamycin, as well as with topical (dermal) formulations of clindamycin. Consider this diagnosis in patients present with diarrhea subsequent to the administration of clindamycin vaginal suppository, because approximately 30% of the clindamycin dose is systemically absorbed from the vagina. Mild cases of pseudomembranous colitis respond to discontinuation of the drug. In moderate to severe cases : management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial drug **PREGNANCY & LACTATION:** Clintas CL suppository should be used during pregnancy only if the potential benefit justifies the potential risk of the fetus. It should be used in lactating women, only if clearly needed. **Interaction:** Intercourse or use of other vaginal products (such as tampons or douches) during treatment with this product should be avoided. These suppositories use an oleaginous base that may weaken latex or rubber products such as condoms or vaginal contraceptive diaphragms. Therefore, the use of such products within 72 hours following treatment with clindamycin vaginal suppository is not recommended. Clindamycin has been shown to have neuromuscular blocking properties that may enhance the action of other neuromuscular blocking agents. when used together, clotrimazole may cause damage to latex contraceptives Patients should be advised to use alternative precautions for at least five days after using clotrimazole **ADVERSE REACTIONS:**Clindamycin : Adverse events reported by 1% or more of patients : Vulvovaginal disorder, Vaginal pain, vaginal monilliasis and fungal infection. Clotrimazole : Rarely patients may experience Local mild burning or irritation immediately after inserting the suppository. Other adverse events : allergic reaction (hypotension, dyspnea, and gastrointestinal disorders), pain, pruritis and rash. (For details, please refer full prescribing information)

(For the use of a registered medical practitioner or hospital or laboratory only)