

AMFY V

COMPOSITION -Lipid based amphotericin B gel 0.1% w/w.**INDICATION:** It is indicated for the treatment of cutaneous and mucocutaneous fungal infections caused by candida (Monilia) species. **DOSAGE & ADMINISTRATION:** Apply Amfy V to the affected areas once or twice daily or as directed by physician.**CONTRAINDICATIONS:** Amfy V should not be applied to patients who have documented hypersensitivity to any of the constituents of this gel.**WARNINGS & PRECAUTIONS:** If there is severe skin rash, redness, burning or itching where the Amfy V is applied, resulting in the worsening of infected lesion, physician must be contacted. Amfy V should not be used if the patient is allergic to amphotericin B. **PREGNANCY AND LACTATION:** Amfy V should be used during pregnancy with caution only if the potential benefit to the mother outweighs the potential risk to the fetus **Interaction:** Amfy V can be used as part of combination with topical or systemic therapy but one should remember that two topical preparations should not be used concomitantly. Other skin medication may affect the effectiveness of amphotericin B.**ADVERSE REACTIONS:** Amfy V is well tolerated. The physician should monitor the patients for any type of adverse event associated with treatment with Amfy V. Avoid using other topical agents at the same time unless prescribed by physician. **OVERDOSE:** An overdose of Amfy V by topical delivery is not likely to occur. If much larger than normal dose has been used and patient does not feel normal, then physician should be contacted. (For details, please refer full prescribing information)

(For the use of a registered medical practitioner or hospital or laboratory only)